

Comment on “Canonical and Microcanonical Calculations for Fermi Systems”

In the context of nuclear physics Pratt [1] recently investigated noninteracting Fermi systems described by the microcanonical and canonical ensemble. As will be shown his discussion of the model of equally spaced levels contains a flaw and a statement which is at least confusing.

In his Letter Pratt gives an alternative derivation of an expression (Eq. (3) of Ref. [1]) in which the canonical partition function $Z_A(\beta)$ of a system with A fermions at inverse temperature β is expressed in terms of the partition function with fewer particles $Z_{A-n}(\beta)$ [2]. Starting with $Z_0(\beta) \equiv 1$, $Z_A(\beta)$ can thus be calculated recursively. This result also leads to an expression in which the canonical occupation probability $\langle n_k \rangle_c$ for a one-particle state with energy ϵ_k is expressed as a sum over terms which contain $Z_{A-n}(\beta)$ and $\exp(-n\beta\epsilon_k)$ [3,1]. $Z_A(\beta)$ can be expressed as the (discrete) Laplace transform of the number of states N_{A,E_r} of a system with A particles and fixed energy E_r . Here the index r indicates, that for a finite system the energy is discrete and given as the sum over A one-particle energies ϵ_k . Furthermore the product $Z_A(\beta) \langle n_k \rangle_c$ is given as the Laplace transform of $N_{A,E_r} \langle n_k \rangle_{mc}$ where $\langle n_k \rangle_{mc}$ is the microcanonical distribution function. Starting with the expressions for the canonical ensemble and using inverse Laplace transformation it is thus possible to obtain recursive relations for N_{A,E_r} [3] and $\langle n_k \rangle_{mc}$. They are given in Eqs. (7) and (8) of Ref. [1].

As an example Pratt investigated a model with equally spaced (spacing g) nondegenerate fermionic levels. The A -particle groundstate energy is given by $E_0(A) = gA(A+1)/2$ and the total energy $E_r(A)$ can be written as $E_r(A) = E_0(A) + gr$, with $r = 0, 1, 2, \dots$. His discussion suggests that there is a direct connection between the number of states with an excitation energy gr and the number of partitions of the integer r into *distinct* parts. This is not the case. Instead the model can be related to the *unrestricted partition problem* [4,5], i.e. the problem of finding the number of ways the integer r can be written as a sum of smaller integers, where it is allowed to *repeatedly use the same integer*. The very fact that the same integer can be used more than once is essential in connection with the mapping of the fermionic system on a bosonic one, a procedure which is known as *bosonization* [4]. For $r \leq A$, N_{A,E_r} is given by the number of partitions $p(r)$ of r which is independent of A . For $r > A$ this mapping breaks down and $N_{A,E_r} < p(r)$ [4,5]. In Fig. 3 of his Letter Pratt presents data for $\langle n_k \rangle_{mc}$ and $r = 20$ calculated using his Eqs. (7) and (8). He does not give the number of particles for which the data have been obtained. We have been informed by Pratt that $r \leq A$, which implies that the data have to be independent of A . For $r \leq A$ the microcanonical occupation probability fulfills the *symmetry relation* $\langle n_k \rangle_{mc} + \langle n_{A-k+1} \rangle_{mc} = 1$ and is a *monotonic* function of k [4,5]. These observations are

essential in connection with the question how the grand canonical distribution function (the Fermi function) is approached in the thermodynamic limit. Both requirements are apparently violated by the data presented by Pratt. We have been informed by the author that in Fig. 3 he accidentally used the wrong set of data. The data shown have been obtained for a model of randomly placed levels. Fig. 1 shows the correct microcanonical occupation probability for the equally spaced level model, $r = 20$, and different system sizes. To obtain the data for $r \leq A$ we have used the three different algorithms presented in Refs. [4], [5], and [1] which all lead to the same curve. For $r > A$ the formulas presented in Refs. [4] and [5] *cannot* be used and only the new result of Pratt which does not make direct use of the mapping to the partition problem is applicable. $\langle n_k \rangle_{mc}$ displays monotonic behavior also for $r > A$. As a test of the validity of our data we have always checked the sum rule $\sum_k \langle n_k \rangle_{mc} = A$ which is fulfilled up to very high precision.




FIG. 1. The microcanonical occupation probability $\langle n_k \rangle_{mc}$ for excitation energy $20g$ and different system sizes. For comparison the Fermi function for temperature $\tau \equiv k_B T/g = 3.49$ is shown as the solid line (see Ref. [1]).

V. Meden and K. Schönhammer
Institut für Theoretische Physik
Universität Göttingen
Bunsenstr. 9
D-37073 Göttingen
Germany

-
- [1] S. Pratt, Phys. Rev. Lett. **84**, 4256 (2000).
 - [2] P. Borrmann and G. Franke, J. Chem. Phys. **98**, 2484 (1993).
 - [3] P. Borrmann *et al.*, Phys. Rev. A **60**, 1519 (1999).
 - [4] K. Schönhammer and V. Meden, Am. J. Phys. **64**, 1168 (1996) and references therein.
 - [5] J. Arnaud *et al.*, Am. J. Phys. **67**, 215 (1999).